

On the Applicability of Post's Lattice[☆]

Michael Thomas

TWT GmbH
Bernhäuser Straße 40–42
73765 Neuhausen auf den Fildern

Abstract

For decision problems $\Pi(B)$ defined over Boolean circuits using gates from a restricted set B only, we have $\Pi(B) \leq_m^{\text{AC}^0} \Pi(B')$ for all finite sets B and B' of gates such that all gates from B can be computed by circuits over gates from B' . In this note, we show that a weaker version of this statement holds for decision problems defined over Boolean formulae, namely that $\Pi(B) \leq_m^{\text{NC}^2} \Pi(B' \cup \{\wedge, \vee\})$ and $\Pi(B) \leq_m^{\text{NC}^2} \Pi(B' \cup \{0, 1\})$ for all finite sets B and B' of Boolean functions such that all $f \in B$ can be defined in B' .

Keywords: computational complexity, Post's lattice

1. Introduction

Let Π denote some decision problem defined over Boolean circuits such that membership in Π is invariant under the substitution of equivalent circuits. Denote by $\Pi(B)$ its restriction to circuits using gates from a finite set B only. It is easily observed that then $\Pi(B) \leq_m^{\text{AC}^0} \Pi(B')$ for all finite sets B and B' such that all gates from B can be computed by circuits over gates from B' (see, e.g., [7, 20]). If we consider formulae instead, this reduction does not necessarily hold; the size of the smallest formula over the Boolean connectives from B' computing some function from B might be of exponential size.

Building on works of [8, 9, 22], we show that a weaker form of this property holds for decision problems defined over formulae, namely that $\Pi(B) \leq_m^{\text{NC}^2} \Pi(B' \cup \{\wedge, \vee\})$ and $\Pi(B) \leq_m^{\text{NC}^2} \Pi(B' \cup \{0, 1\})$ for all finite sets B and B' of Boolean functions such that all $f \in B$ can be defined in B' . Moreover, if all connectives in B can be expressed using either only conjunction (\wedge), only disjunction (\vee) or only the exclusive-or (\oplus), we obtain $\Pi(B) \leq_m^{\text{AC}^0} \Pi(B')$, as in the circuit setting.

These results provide a (partial) account for the polytomous complexity classifications of problems parametrized by the set of available Boolean connectives: the complexity of the satisfiability problem was, for instance, shown to be NP-complete if $x \leftrightarrow y$ can be composed from the available Boolean connectives, and solvable in logspace in all other cases [13]. Further results include a variety of problems in propositional logic [5, 19], modal logics [1], temporal logics [2–4, 16], their hybrid variants [14, 15], and nonmonotonic logics [6, 11, 23].

We point out that the results obtained herein are completely general in that they do not rely on properties of the considered

problems except invariance of membership under substitution of logically equivalent formulae (i.e., if (φ, x) is an instance of Π with φ being a Boolean formula and if φ' is a Boolean formula logically equivalent to φ , then $(\varphi, x) \in \Pi$ iff $(\varphi', x) \in \Pi$). This generality comes at the price of a fairly powerful reduction. However, in practice, most problems exhibit additional structure that allow to further restrict the notion of reductions considered.

2. Preliminaries

Propositional Logic. Let \mathcal{L} be the set of propositional formulae, i.e., the set of formulae defined via

$$\varphi ::= a \mid c(\varphi, \dots, \varphi),$$

where a is a proposition and c is an n -ary connective. We associate an n -ary connective c with the n -ary Boolean function $f_c: \{0, 1\}^n \rightarrow \{0, 1\}$ defined by $f_c(a_1, \dots, a_n) := 1$ if and only if the formula $c(x_1, \dots, x_n)$ becomes true when assigning a_i to c_i , $1 \leq i \leq n$. Let $\varphi_{[\alpha/\beta]}$ denote φ with all occurrences of the subformula α replaced by some formula β . For a finite set B of Boolean connectives, let $\mathcal{L}(B)$ denote the set of B -formulae, i.e., the set \mathcal{L} restricted to formulae using connectives from B only. The depth of a formula is the maximum nesting depth of Boolean connectives; the size of a formula is equal to the number of symbols used to represent it.

Clones and Post's Lattice. A clone is a set of Boolean functions that is closed under superposition, i.e., B contains all projections (the functions $f(x_1, \dots, x_n) = x_k$ for all $1 \leq k \leq n$) and is closed under arbitrary composition [17]. For a set B of Boolean functions, we denote by $[B]$ the smallest clone containing B and call B a base for $[B]$. A B -formula g is called B -representation of f if f and g are equivalent, i.e., $f \equiv g$. It is clear that B -representations exist for every $f \in [B]$.

[☆]Work supported by DFG grant VO 630/6-2 and performed while employed at the Gottfried Wilhelm Leibniz Universität Hannover.

Email address: michael.thomas@twt-gmbh.de (or thomas@thi.uni-hanover.de)

In [18], Post showed that the set of all clones ordered by inclusion together with $[A \cap B]$ and $[A \cup B]$ forms a lattice and found a finite base for each clone, see Figure 1. To introduce the clones, we define the following properties. Say that a set $A \subseteq \{0, 1\}^n$ is c -separating, $c \in \{0, 1\}$, if there exists an $i \in \{1, \dots, n\}$ such that $(a_1, \dots, a_n) \in A$ implies $a_i = c$. Let f be an n -ary Boolean function and define the dual of f to be the Boolean function $\text{dual}(f)(x_1, \dots, x_n) := \neg f(\neg x_1, \dots, \neg x_n)$. We say that

- f is c -reproducing if $f(c, \dots, c) = c$, $c \in \{0, 1\}$;
- f is c -separating if $f^{-1}(c)$ is c -separating, $c \in \{0, 1\}$;
- f is c -separating of degree m if all $A \subseteq f^{-1}(c)$ with $|A| = m$ are c -separating;
- f is monotone if $a_1 \leq b_1, a_2 \leq b_2, \dots, a_n \leq b_n$ implies $f(a_1, \dots, a_n) \leq f(b_1, \dots, b_n)$;
- f is self-dual if $f \equiv \text{dual}(f)$;
- f is affine if $f(x_1, \dots, x_n) \equiv x_1 \oplus \dots \oplus x_n \oplus c$ with $c \in \{0, 1\}$;
- f is essentially unary if f depends on at most one variable.

The above properties canonically extend to sets B of Boolean functions by requiring that all $f \in B$ satisfy the given property. The list of all clones is given in Table 1.

Reductions. Let A and B be decision problems. Say that A C -many-one reduces to B (written: $A \leq_m^C B$) if there exists a C -computable function f mapping instances x of A to instances $f(x)$ of B such that $x \in A \iff f(x) \in B$. If $A \leq_m^C B$ and $B \leq_m^C A$, we also write $A \equiv_m^C B$.

3. Previous Results and Auxiliary Lemmas

The following lemma due to Spira is well-known and will be useful if the given set of Boolean functions is functionally complete.

Lemma 3.1 ([22]) *Let φ be a propositional formula. Then there exists an equivalent $\{\wedge, \vee, \neg\}$ -formula ψ such that the depth of ψ is $O(\log |\varphi|)$ and the size of ψ is $|\varphi|^{O(1)}$.*

Lemma 3.2 *Let φ be a propositional formula over Boolean connectives from $[B] \subseteq \mathbb{M}$ and let $g(x, y, z) := x \vee (y \wedge z)$. Then there exists an equivalent $(B \cup \{g, 0, 1\})$ -formula ψ such that the depth of ψ is $O(\log |\varphi|)$ and the size of ψ is $|\varphi|^{O(1)}$.*

Proof. We proceed analogous to a construction of Bonnet and Buss from [8]. Let φ be the given formula over connectives from a set B and let m be the number of occurrences of propositions in φ . We claim that there exists an equivalent $(B \cup \{g, 0, 1\})$ -formula of depth $O(\log m)$ and polynomial size.

If $m \leq 1$ then φ is equivalent to x or a constant and can be implemented in depth 1. Hence assume that $m > 1$ and that the claim holds for all smaller m . Then there exists a subformula ψ that contains $\geq \frac{m}{k}$ occurrences of propositions, where k is a bound on the arity of the functions in B (see also [9]). Define

$\varphi' := g(\varphi_{[\psi/0]}, \varphi_{[\psi/1]}, \psi) \equiv \varphi_{[\psi/0]} \vee (\varphi_{[\psi/1]} \wedge \psi)$. By monotonicity, φ is equivalent to φ' . Moreover, by induction hypothesis, we may assume the depths of ψ and $\varphi_{[\psi/c]}$, $c \in \{0, 1\}$, to be $O(\log \frac{m}{k})$ and $O(\log \frac{(k-1)m}{k})$, respectively. Denote by d the constant hidden in these O -notations. Then the depth of φ' can be bounded by $2 + \max\{\text{depth}(\varphi_{[\psi/0]}), \text{depth}(\varphi_{[\psi/1]}), \text{depth}(\psi)\} = 2 + d \cdot k \cdot \log \left(\frac{(k-1)m}{k}\right) = 2 + d \cdot k \cdot \left(\log m + \log\left(1 - \frac{1}{k}\right)\right) < 2 + d \cdot k \cdot \left(\log m - \frac{1}{k}\right) \in O(\log m)$, as $\log\left(1 - \frac{1}{k}\right) < -\frac{1}{k}$. Concluding, the size of φ' is at most quadratic in the size of φ . \square

Lemma 3.3 *Let φ be a propositional formula over Boolean connectives from $B \subseteq \mathbb{M}$ and let $h(x, y, z) := x \wedge (y \vee z)$. Then there exists an equivalent $(B \cup \{h, 0, 1\})$ -formula ψ such that the depth of ψ is $O(\log |\varphi|)$ and the size of ψ is $|\varphi|^{O(1)}$.*

Proof. Analogous to Lemma 3.2 using $\varphi' := h(\varphi_{[\psi/1]}, \varphi_{[\psi/0]}, \psi) \equiv \varphi_{[\psi/1]} \wedge (\varphi_{[\psi/0]} \vee \psi)$ in the inductive step. \square

4. Results

Throughout this section, let B and B' be for finite sets of Boolean connectives and Σ be an alphabet. We will first formalize the notion of problems defined over propositional formulae and invariance under the substitution of equivalent B -formulae.

Definition 4.1 *A decision problem defined over (propositional) formulae is any set of $\Pi \subseteq \Sigma^* \times \mathcal{L}$. We will write $\Pi(B)$ for $\Pi \cap (\Sigma^* \times \mathcal{L}(B))$.*

Further, say that a decision problem $\Pi(B)$ defined over propositional formulae is invariant under the substitution of equivalent formulae if $(\varphi, x) \in \Pi$ if and only if $(\psi, x) \in \Pi$ for all formulae ψ equivalent to φ .

Lemma 4.2 *Fix B and let $\Pi(B)$ be a decision problem defined over propositional formulae that is invariant under the substitution of equivalent formulae. Then the following holds for all B' satisfying $B \subseteq [B']$:*

1. *If $[B] \subseteq \mathbb{E}$ or $[B] \subseteq \mathbb{V}$, then $\Pi(B) \leq_m^{\text{AC}^0} \Pi(B')$.*
2. *If $[B] \subseteq \mathbb{L}$, then $\Pi(B) \leq_m^{\text{AC}^0[2]} \Pi(B')$.*

Proof. First suppose that $[B] \subseteq \mathbb{E}$ and let $\Pi(B)$ be as in the statement of the lemma. Then any B -formula φ over propositions x_1, x_2, \dots is equivalent to a formula $\varphi' := c \wedge \bigwedge_{i \in I} x_i$, where $c \in \{0, 1\}$. This representation is computable in logarithmic space, as $c = 0$ iff φ is not satisfied by the assignment setting all propositions to 1 (i.e., $\varphi(1, \dots, 1) = 0$), and $i \in I$ iff $\varphi(1, \dots, 1) = 1$ and φ is not satisfied by the assignment setting all propositions but x_i to 1. By inserting parentheses, φ' can be transformed into a formula of logarithmic depth such that replacing all occurring constants and connectives with their B' -representations yields an equivalent B' -formula φ'' of size at most $2^{O(\log |\varphi|)} = |\varphi|^{O(1)}$. Thus, given input $(\varphi, x) \in \mathcal{L}(B) \times \Sigma^*$, it suffices to construct (φ', x) . As the evaluation of B -formulae for $[B] \subseteq \mathbb{E}$ can be performed in AC^0 [21], we finally obtain $\Pi(B) \leq_m^{\text{AC}^0} \Pi(B')$ for all B' satisfying $B \subseteq [B']$.

For $[B] \subseteq \mathbb{L}$ and $[B] \subseteq \mathbb{V}$, similar arguments work. The construction of $\varphi' := c \oplus \bigoplus_{i \in I} x_i$ (resp. $\varphi' := c \vee \bigvee_{i \in I} x_i$) is as

Clone	Definition	Base
BF	All Boolean functions	$\{x \wedge y, \neg x\}$
R ₀	$\{f \in \text{BF} \mid f \text{ is 0-reproducing}\}$	$\{x \wedge y, x \oplus y\}$
R ₁	$\{f \in \text{BF} \mid f \text{ is 1-reproducing}\}$	$\{x \vee y, x \leftrightarrow y\}$
R ₂	$R_0 \cap R_1$	$\{x \vee y, x \wedge (y \leftrightarrow z)\}$
M	$\{f \in \text{BF} \mid f \text{ is monotone}\}$	$\{x \wedge y, x \vee y, 0, 1\}$
M ₀	$M \cap R_0$	$\{x \wedge y, x \vee y, 0\}$
M ₁	$M \cap R_1$	$\{x \wedge y, x \vee y, 1\}$
M ₂	$M \cap R_2$	$\{x \wedge y, x \vee y\}$
S ₀	$\{f \in \text{BF} \mid f \text{ is 0-separating}\}$	$\{x \rightarrow y\}$
S ₀ ⁿ	$\{f \in \text{BF} \mid f \text{ is 0-separating of degree } n\}$	$\{x \rightarrow y, \text{dual}(t_n^{n+1})\}$
S ₁	$\{f \in \text{BF} \mid f \text{ is 1-separating}\}$	$\{x \leftrightarrow y\}$
S ₁ ⁿ	$\{f \in \text{BF} \mid f \text{ is 1-separating of degree } n\}$	$\{x \leftrightarrow y, t_n^{n+1}\}$
S ₀₂ ⁿ	$S_0^n \cap R_2$	$\{x \vee (y \wedge \neg z), \text{dual}(t_n^{n+1})\}$
S ₀₂	$S_0 \cap R_2$	$\{x \vee (y \wedge \neg z)\}$
S ₀₁ ⁿ	$S_0^n \cap M$	$\{\text{dual}(t_n^{n+1}), 1\}$
S ₀₁	$S_0 \cap M$	$\{x \vee (y \wedge z), 1\}$
S ₀₀ ⁿ	$S_0^n \cap R_2 \cap M$	$\{x \vee (y \wedge z), \text{dual}(t_n^{n+1})\}$
S ₀₀	$S_0 \cap R_2 \cap M$	$\{x \vee (y \wedge z)\}$
S ₁₂ ⁿ	$S_1^n \cap R_2$	$\{x \wedge (y \vee \neg z), t_n^{n+1}\}$
S ₁₂	$S_1 \cap R_2$	$\{x \wedge (y \vee \neg z)\}$
S ₁₁ ⁿ	$S_1^n \cap M$	$\{t_n^{n+1}, 0\}$
S ₁₁	$S_1 \cap M$	$\{x \wedge (y \vee z), 0\}$
S ₁₀ ⁿ	$S_1^n \cap R_2 \cap M$	$\{x \wedge (y \vee z), t_n^{n+1}\}$
S ₁₀	$S_1 \cap R_2 \cap M$	$\{x \wedge (y \vee z)\}$
D	$\{f \in \text{BF} \mid f \text{ is self-dual}\}$	$\{(x \wedge y) \vee (x \wedge \neg z) \vee (\neg y \wedge \neg z)\}$
D ₁	$D \cap R_2$	$\{(x \wedge y) \vee (x \wedge \neg z) \vee (y \wedge \neg z)\}$
D ₂	$D \cap M$	$\{(x \wedge y) \vee (x \wedge z) \vee (y \wedge z)\}$
L	$\{f \in \text{BF} \mid f \text{ is affine}\}$	$\{x \oplus y, 1\}$
L ₀	$L \cap R_0$	$\{x \oplus y\}$
L ₁	$L \cap R_1$	$\{x \leftrightarrow y\}$
L ₂	$L \cap R_2$	$\{x \oplus y \oplus z\}$
L ₃	$L \cap D$	$\{x \oplus y \oplus z \oplus 1\}$
E	$\{f \in \text{BF} \mid f \text{ is constant or a conjunction}\}$	$\{x \wedge y, 0, 1\}$
E ₀	$E \cap R_0$	$\{x \wedge y, 0\}$
E ₁	$E \cap R_1$	$\{x \wedge y, 1\}$
E ₂	$E \cap R_2$	$\{x \wedge y\}$
V	$\{f \in \text{BF} \mid f \text{ is constant or a disjunction}\}$	$\{x \vee y, 0, 1\}$
V ₀	$V \cap R_0$	$\{x \vee y, 0\}$
V ₁	$V \cap R_1$	$\{x \vee y, 1\}$
V ₂	$V \cap R_2$	$\{x \vee y\}$
N	$\{f \in \text{BF} \mid f \text{ is essentially unary}\}$	$\{\neg x, 0, 1\}$
N ₂	$N \cap D$	$\{\neg x\}$
I	$\{f \in \text{BF} \mid f \text{ is constant or a projection}\}$	$\{\text{id}, 0, 1\}$
I ₀	$I \cap R_0$	$\{\text{id}, 0\}$
I ₁	$I \cap R_1$	$\{\text{id}, 1\}$
I ₂	$I \cap R_2$	$\{\text{id}\}$

Table 1: List of all clones with definition and bases, where id denotes the identity and $t_n^{n+1}(x_0, \dots, x_n) := \bigvee_{i=0}^n (x_0 \wedge \dots \wedge x_{i-1} \wedge x_{i+1} \wedge \dots \wedge x_n)$.

Figure 1: Post's lattice

follows: $c \equiv 1$ iff $\varphi(0, \dots, 0) = 1$, and $i \in I$ iff the truth value of φ under the assignment setting all propositions to 0 and the truth value of φ under the assignment setting only the proposition x_i to 1 differ (resp. $i \in I$ iff $\varphi(0, \dots, 0) = 0$ and φ is satisfied by the assignment setting only the proposition x_i to 1). And the evaluation of B -formulae for $[B] \subseteq V$ can be performed in AC^0 , while for $[B] \subseteq L$ we require $AC^0[2]$. \square

Henceforth, let $C \supseteq AC^0$ be such that given φ the formula ψ in the Lemmas 3.1, 3.2 and 3.3 can be computed in C . (A direct implementation of these restructurings requires $O(\log^2 n)$ space, hence $NC^2 \subseteq C$ suffices; Cook and Gupta showed that Spira's construction can actually be performed in alternating $O(\log n \cdot \log \log n)$ -time [12]).

Lemma 4.3 *Fix B and let $\Pi(B)$ be a decision problem defined over propositional formulae that is invariant under the substitution of equivalent formulae. Then the following holds for all B' satisfying $B \subseteq [B']$:*

1. If $S_{00} \subseteq [B] \subseteq M$, then $\Pi(B) \leq_m^C \Pi(B' \cup \{\wedge\})$.
2. If $S_{10} \subseteq [B] \subseteq M$, then $\Pi(B) \leq_m^C \Pi(B' \cup \{\vee\})$.

Proof. Suppose that $S_{00} \subseteq [B] \subseteq M$ and let $\Pi(B)$ be as in the statement of the lemma. Let (φ, x) be the given instance with $\varphi \in \mathcal{L}(B)$. Denote by $g(x, y, z)$ the function $x \vee (y \wedge z) \in S_{00} \subseteq [B]$. Then, by Lemma 3.2, there exists a $(B \cup \{g, 0, 1\})$ -formula φ' of logarithmic depth and polynomial size such that $\varphi \equiv \varphi'$. Obtain φ' from φ by replacing all connectives from $B \cup \{g\}$ with their B' -representations. Next, if $1 \notin [B']$, we eliminate the constant 1 by replacing it with the B' -representation of $\bigvee_{i=1}^n x_i$, where x_1, \dots, x_n enumerate all propositions occurring in φ' . Analogously, if $0 \notin [B']$, we eliminate the constant 0 by replacing it with the B' -representation of $\bigwedge_{i=1}^n x_i$. Call the resulting formula φ'' . If $1 \notin [B']$, then φ cannot be satisfied by the assignment setting all propositions to 0, as $[B'] \subseteq R_0$; for all other assignments, $\bigvee_{i=1}^n x_i$ is satisfied. If $0 \notin [B']$, then φ is satisfied by the assignment setting all propositions to 1, as $[B] \subseteq R_1$; for all other assignments, $\bigwedge_{i=1}^n x_i$ is not satisfied. Therefore, φ'' is equivalent to φ .

Consequently, the mapping $(\varphi, x) \mapsto (\varphi'', x)$ constitutes a \leq_m^C -reduction from $\Pi(B)$ to $\Pi(B' \cup \{\wedge\})$, as φ' is C -computable by assumption and the construction of φ'' from φ' requires local replacements only. This concludes the proof of the first claim.

As for the second claim, suppose that $S_{10} \subseteq [B] \subseteq M$. Denote again by (φ, x) the given instance and abbreviate with $h(x, y, z)$ the function $x \wedge (y \vee z) \in S_{10} \subseteq [B]$. By Lemma 3.3, there exists a $(B \cup \{h, 0, 1\})$ -formula φ' of logarithmic depth and polynomial size such that $\varphi \equiv \varphi'$. Obtain φ'' from φ by replacing all connectives from $B \cup \{h\}$ with their B' -representations and eliminating the constants not contained in $[B']$ as above. Then $(\varphi, x) \mapsto (\varphi'', x)$ constitutes a \leq_m^C -reduction from $\Pi(B)$ to $\Pi(B' \cup \{\vee\})$. \square

Lemma 4.4 *Fix B and let $\Pi(B)$ be a decision problem defined over propositional formulae that is invariant under the substitution of equivalent formulae. Then the following holds for all B' satisfying $B \subseteq [B']$:*

1. If $S_{02} \subseteq [B]$, then $\Pi(B) \leq_m^C \Pi(B' \cup \{\wedge\})$.
2. If $S_{12} \subseteq [B]$, then $\Pi(B) \leq_m^C \Pi(B' \cup \{\vee\})$.

Proof. Suppose that $S_{02} \subseteq [B]$ and let $\Pi(B)$ be as in the statement of the lemma. Let (φ, x) be the given instance with $\varphi \in \mathcal{L}(B)$. By Lemma 3.1, there exists a $\{\wedge, \vee, \neg\}$ -formula φ' of logarithmic depth and polynomial size such that $\varphi \equiv \varphi'$. Observe that φ' can be constructed from φ by a procedure similar to that used in the proof of Lemma 3.2 (in the inductive step, use $(\varphi_{[\psi/0]} \wedge \neg\psi) \vee (\varphi_{[\psi/1]} \wedge \psi)$ as the new formula). As $x \vee (y \wedge \neg z)$ is a base for $[B']$ and $x \vee (y \wedge \neg 0) \equiv x \vee y, 0 \vee (y \wedge \neg(0 \vee (1 \wedge \neg z))) \equiv y \wedge z$ and $0 \vee (1 \wedge \neg z) \equiv \neg z$, we obtain $\{\wedge, \vee, \neg\} \in [B' \cup \{0, 1\}]$. So we can first replace all connectives from $B \cup \{\wedge, \vee, \neg\}$ in φ' with their $(B' \cup \{0, 1\})$ -representations, and second, eliminate those constants not contained in $[B']$ as in the proof of Lemma 4.3. Call the resulting formula φ'' . As $S_{02} \subseteq [B']$ and $1 \notin [B']$ imply that $[B'] \subseteq R_0$, and $S_{02} \subseteq [B']$ and $0 \notin [B']$ imply that $[B'] \subseteq R_1$, φ'' is equivalent to φ' by the same arguments as above. The function mapping (φ, x) to (φ'', x) is hence a \leq_m^C -reduction from $\Pi(B)$ to $\Pi(B' \cup \{\wedge\})$.

The proof of the second claim is analogous. \square

Lemma 4.5 *Fix B and let $\Pi(B)$ be a decision problem defined over propositional formulae that is invariant under the substitution of equivalent formulae. If $D_2 \subseteq [B] \subseteq D$, then $\Pi(B) \leq_m^C \Pi(B' \cup \{\vee\})$ and $\Pi(B) \leq_m^C \Pi(B' \cup \{\wedge\})$ for all B' satisfying $B \subseteq [B']$.*

Proof. Let B and $\Pi(B)$ be as in the statement of the lemma and denote by (φ, x) the given instance with $\varphi \in \mathcal{L}(B)$. On the one hand, if $[B] = D_2$, then by Lemma 3.2 and Lemma 3.3 there exist logarithmic-depth polynomial-size formulae $\varphi' \in \mathcal{L}(B \cup \{x \vee (y \wedge z), 0, 1\})$ and $\varphi'' \in \mathcal{L}(B \cup \{x \wedge (y \vee z), 0, 1\})$. Proceeding as in the proof of Lemma 4.3, we obtain the desired reduction.

On the other hand, if $D_1 \subseteq [B]$, then by Lemma 3.1 there exists a $\{\wedge, \vee, \neg\}$ -formula φ' of logarithmic depth and polynomial size such that $\varphi \equiv \varphi'$. As $[B' \cup \{0, 1\}] = BF$, we may replace all connectives in φ' with their $(B' \cup \{0, 1\})$ -representations. If $[B'] \subseteq R_0$ (or if $[B'] \subseteq R_1$), we may eliminate the constant 1 (or 0) as in the proof of Lemma 4.3. Otherwise, if $[B'] = BF$, then we may replace 1 with $t \vee \neg t$ and 0 with $t \wedge \neg t$, where t is an arbitrary fresh proposition. Either way, we obtain a formula $\varphi'' \in \mathcal{L}(B' \cup C)$ of polynomial size such that $\varphi'' \equiv \varphi$ and C is either $\{\vee\}$, $\{\wedge\}$, or the empty set. The mapping from $(\varphi, x) \in \Pi(B)$ to $(\varphi'', x) \in \Pi(B' \cup C)$ is the desired \leq_m^C -reduction. \square

We are now ready to state our main theorem.

Theorem 4.6 *Fix B and let $\Pi(B)$ be a decision problem defined over propositional formulae that is invariant under the substitution of equivalent formulae. Then the following holds for all B' satisfying $B \subseteq [B']$:*

- If $[B] \subseteq V$ or $[B] \subseteq L$ or $[B] \subseteq E$ or $M_2 \subseteq [B]$, then $\Pi(B) \leq_m^C \Pi(B')$.
- If $S_{00} \subseteq [B] \subseteq S_0^2$ or $D_2 \subseteq [B] \subseteq D$, then $\Pi(B) \leq_m^C \Pi(B' \cup \{\wedge\})$.

- If $S_{10} \subseteq [B] \subseteq S_1^2$ or $D_2 \subseteq [B] \subseteq D$, then $\Pi(B) \leq_m^C \Pi(B' \cup \{\vee\})$.

Proof. Consider the lattice in Fig. 1. It holds that either (a) $[B] \subseteq V$, (b) $[B] \subseteq L$, (c) $[B] \subseteq E$, (d) $S_{00} \subseteq [B] \subseteq S_0^2$, (e) $S_{10} \subseteq [B] \subseteq S_1^2$, (f) $D_2 \subseteq [B] \subseteq D$, or (g) $M_2 \subseteq [B]$. The first claim corresponds to the cases (a)–(c) and (g). The second and third claim correspond to case (d) and (f) resp. (e) and (f).

In cases (a)–(c), $\Pi(B) \leq_m^C \Pi(B')$ follows from Lemma 4.2.

As for case (d), we have either $[B] \subseteq S_{01}^2$ or $S_{02} \subseteq [B]$. In either case, the reduction $\Pi(B) \leq_m^C \Pi(B' \cup \{\wedge\})$ is implied by Lemmas 4.3 and 4.4.

Case (e) analogously yields $\Pi(B) \leq_m^C \Pi(B' \cup \{\vee\})$.

For case (f), Lemma 4.5 yields both $\Pi(B) \leq_m^C \Pi(B' \cup \{\wedge\})$ and $\Pi(B) \leq_m^C \Pi(B' \cup \{\vee\})$.

It remains to consider case (g): Fix a set B with $M_2 \subseteq [B]$. If we suppose that $[B] \subseteq M$, Lemma 4.3 yields $\Pi(B) \leq_m^C \Pi(B' \cup \{\wedge\})$ (or $\Pi(B) \leq_m^C \Pi(B' \cup \{\vee\})$). Yet, for all such B , we have $\{\wedge, \vee\} \in [B] \subseteq [B']$; in which case Lemma 4.3 actually yields $\Pi(B) \leq_m^C \Pi(B')$. The same argument applies if $M \not\subseteq [B]$, using Lemma 4.4 instead. This completes the last case and establishes the theorem. \square

As an easy consequence of Theorem 4.6 and the remark below Lemma 4.2, we obtain the following two corollaries:

Corollary 4.7 *If $\Pi(B \cup \{0, 1\}) \leq_m^C \Pi(B)$ for all B , then $\Pi(B) \leq_m^C \Pi(B')$ for all B and B' such that $B \subseteq [B']$; in particular, $\Pi(B)$ is C -equivalent to Π restricted to one of the following sets of functions: $\{\wedge, \vee, \neg\}$, $\{\wedge, \vee\}$, $\{\wedge\}$, $\{\vee\}$, $\{\oplus\}$, $\{\neg\}$, $\{\text{id}\}$.*

Corollary 4.8 *Let B be a finite set of Boolean functions.*

- If $[B] = \text{BF}$, then $\Pi(B) \equiv_m^{\text{NC}^2} \Pi(\{\wedge, \vee, \neg\})$.
- If $[B] = M$, then $\Pi(B) \equiv_m^{\text{NC}^2} \Pi(\{\wedge, \vee, 0, 1\})$.
- If $[B] = L$, then $\Pi(B) \equiv_m^{\text{AC}^0[2]} \Pi(\{\oplus, 1\})$.
- If $[B] = N$, then $\Pi(B) \equiv_m^{\text{AC}^0[2]} \Pi(\{\neg, 1\})$.
- If $[B] = E$, then $\Pi(B) \equiv_m^{\text{AC}^0} \Pi(\{\wedge, 0, 1\})$.
- If $[B] = V$, then $\Pi(B) \equiv_m^{\text{AC}^0} \Pi(\{\vee, 0, 1\})$.

It is straightforward to extend Corollary 4.8 to those clones not containing both constants.

5. Concluding Remarks

The results presented in this note provide insight into why complexity classifications of problems in Post’s lattice yield only a finite number of complexity degrees.

These results are completely general in the sense that we did not place any restrictions on the considered decision problems Π (unless, of course, that membership in Π is invariant under substitution of equivalent formulae). However, typically instances of natural decision problems exhibit additional structure; by exploiting this structure one may further reduce the computational power of the reduction \leq_m^C , or obtain $\Pi(B) \leq_m^C \Pi(B')$ without resorting to the assumption $\Pi(B \cup \{0, 1\}) \leq_m^C$

$\Pi(B')$ given Corollary 4.7. For example, if $\Pi(\{\wedge, \vee\}) \leq_m^C \Pi(B)$ for all finite sets B of Boolean functions satisfying $S_{00} \subseteq [B]$ or $S_{10} \subseteq [B]$ or $D_2 \subseteq [B]$, then $\Pi(B) \leq_m^C \Pi(B')$ for all B and B' satisfying $B \subseteq [B']$. This holds for the propositional implication problem [5], among others.

It is worth noting that, on the other hand, there exist natural problems that do not satisfy the conditions imposed on Π above. Amongst those is the problem BFMIN, which asks to determine, given a Boolean formula and an integer k , whether there exists an equivalent formula of size $\leq k$. This problem has recently been shown to be Σ_2^P -complete for the Boolean standard base $B = \{\wedge, \vee, \neg\}$ using Turing reductions [10]. However, considering its restriction to B -formulae, we obtain $\text{BFMIN}(B) \not\leq_m^C \text{BFMIN}(B')$: Let $\varphi \notin \mathcal{L}(B)$ be some Boolean formula of arity n . Then $(\varphi, c(n)) \in \text{BFMIN}(B \cup \{\varphi\})$ for some constant c (depending on n only), while $(\varphi, k) \notin \text{BFMIN}(B)$ for all $k \in \mathbb{N}$.

References

- [1] M. Bauland, E. Hemaspaandra, H. Schnoor, and I. Schnoor. Generalized modal satisfiability. In *Proc. 23rd Annual Symposium on Theoretical Aspects of Computer Science*, volume 3884 of *Lecture Notes in Computer Science*, pages 500–511. Springer, 2006.
- [2] M. Bauland, M. Mundhenk, T. Schneider, H. Schnoor, I. Schnoor, and H. Vollmer. The tractability of model-checking for LTL: The good, the bad, and the ugly fragments. *Electr. Notes Theor. Comput. Sci.*, 231:277–292, 2009.
- [3] M. Bauland, T. Schneider, H. Schnoor, I. Schnoor, and H. Vollmer. The complexity of generalized satisfiability for linear temporal logic. *Logical Methods in Computer Science*, 5(1), 2008.
- [4] O. Beyersdorff, A. Meier, M. Mundhenk, T. Schneider, M. Thomas, and H. Vollmer. Model checking CTL is almost always inherently sequential. In *Proc. 16th International Symposium on Temporal Representation and Reasoning*, pages 21–28. IEEE Computer Society, 2009.
- [5] O. Beyersdorff, A. Meier, M. Thomas, and H. Vollmer. The complexity of propositional implication. *Inf. Process. Lett.*, 109(18):1071–1077, 2009.
- [6] O. Beyersdorff, A. Meier, M. Thomas, and H. Vollmer. The complexity of reasoning for fragments of default logic. In *Proc. 12th International Conference on Theory and Applications of Satisfiability Testing*, volume 5584 of *Lecture Notes in Computer Science*, pages 51–64. Springer, 2009.
- [7] E. Böhler, N. Creignou, S. Reith, and H. Vollmer. Playing with Boolean blocks, part I: Post’s lattice with applications to complexity theory. *ACM-SIGACT Newsletter*, 34(4):38–52, 2003.
- [8] M. L. Bonet and S. R. Buss. Size-depth tradeoffs for Boolean formulae. *Information Processing Letters*, 49(3):151–155, 1994.
- [9] R. P. Brent. The parallel evaluation of general arithmetic expressions. *J. ACM*, 21(2):201–206, 1974.
- [10] D. Buchfuhrer and C. Umans. The complexity of Boolean formula minimization. *J. Comput. Syst. Sci.*, 77(1):142–153, 2011.
- [11] N. Creignou, A. Meier, M. Thomas, and H. Vollmer. The complexity of reasoning for fragments of autoepistemic logic. In *Circuits, Logic, and Games*, volume 10061 of *Dagstuhl Seminar Proceedings*, 2010.
- [12] A. Gupta. A fast parallel algorithm for recognition of parenthesis languages. Technical Report TR-182/85, Department of Computer Science, University of Toronto, 1985.
- [13] H. R. Lewis. Satisfiability problems for propositional calculi. *Mathematical Systems Theory*, 13:45–53, 1979.
- [14] A. Meier, M. Mundhenk, T. Schneider, M. Thomas, V. Weber, and F. Weiss. The complexity of satisfiability for fragments of hybrid logic—Part I. *Journal of Applied Logic*, 8(4):409–421, 2010.
- [15] A. Meier, M. Mundhenk, T. Schneider, M. Thomas, and F. Weiss. The complexity of satisfiability for fragments of hybrid logic—Part II. In *Proc. International Workshop on Hybrid Logic*, Electronic Notes in Theoretical Computer Science. Elsevier, 2010.
- [16] A. Meier, M. Mundhenk, M. Thomas, and H. Vollmer. The complexity of satisfiability for fragments of CTL and CTL*. *Int. J. Found. Comput. Sci.*, 20(5):901–918, 2009.

- [17] N. Pippenger. *Theories of Computability*. Cambridge University Press, 1997.
- [18] E. Post. The two-valued iterative systems of mathematical logic. *Annals of Mathematical Studies*, 5:1–122, 1941.
- [19] S. Reith. On the complexity of some equivalence problems for propositional calculi. In *Proc. 28th International Symposium on Mathematical Foundations of Computer Science*, volume 2747 of *Lecture Notes in Computer Science*, pages 632–641. Springer, 2003.
- [20] S. Reith and K. W. Wagner. The complexity of problems defined by Boolean circuits. In *Proceedings of the Conference Mathematical Foundation of Informatics 1999*, pages 141–156. World Scientific Publishing, 2000.
- [21] H. Schnoor. The complexity of model checking for Boolean formulas. *Int. J. Found. Comput. Sci.*, 21(3):289–309, 2010.
- [22] P. M. Spira. On time-hardware complexity tradeoffs for Boolean functions. In *Proc. 4th Hawaii International Symposium on System Sciences*, pages 525–527, 1971.
- [23] M. Thomas. The complexity of circumscriptive inference in Post’s lattice. In *Proc. 10th International Conference on Logic Programming and Nonmonotonic Reasoning*, volume 5753 of *Lecture Notes in Computer Science*, pages 290–302. Springer, 2009.